

Dear Patient:

## Abraham Jacob, M.D. Department of Surgery/Division of Otolaryngology Otology, Neurotology, and Cranial Base Surgery

1501 N. Campbell Ave., 5OPC Tucson, AZ 85724 Office: (520) 694-1000 Fax: (520) 694-6101

### BALANCE DISORDERS/ENG/VNG PRE-TEST INSTRUCTIONS

#### IMPORTANT PRE-TEST INSTRUCTIONS FOR INNER EAR BALANCE TESTING

Yo	u have been scheduled for the following tests on	_at	_ a.m./p.m.
Please note, if you arrive more than 15 minutes late you will be asked to reschedule.			
	ENG/VNG (electronystagmography) (testing takes approximately 75 minute	s)	
	Audiogram (hearing test) (testing takes approximately 30 minutes)		
1.	<b>IMPORTANT INFORMATION</b> : Accurate inner ear function testing requires a your central nervous system or that suppress you inner ear function to be <u>signarior to the testing appointment</u> . This would include <u>any</u> medications you including Antivert, Meclizine, Valium, Dramamine, or scopolamine patches. the above medications in the 48 hours prior to your testing appointment, <u>we your test and we will need to reschedule your appointment in order to obtain</u>	e stopped a full 48 hours ou take for dizziness, s. If you forget and take any of we will be unable to perform	

- 2. Other CNS acting medications that need to be stopped 48 hours prior to your appointment would include, but are not limited to: Anti-depressants (Zoloft, Prozac, Wellbutrin), sleeping pills, tranquilizers, anti-anxiety medications, sedatives, prescription pain killers that contain narcotics (Tylenol #3, etc.), any cold or allergy medications that make you sleepy such as Benadryl, Nyquil etc. However, some medications should not be stopped abruptly. Please check with your pharmacist or the physician who wrote the prescription with any questions or concerns regarding stopping these medications. If your physician does not want you to stop any of the meds mentioned in this paragraph, please let the clinic know when you present for your appointment.
- 3. Do not drink alcohol for 48 hours prior to this appointment. (includes hard liquor, wine, or beer)
- 4. IMPORTANT INFORMATION: CONTINUE THESE TYPES OF MEDICATIONS: Continue anything you take for heart or kidney problems, high blood pressure, curculatory disorders, blood thinners, breathing disorders, diabetes, cancer, arthritis (non-narcotics), seizures, or hormone imbalance. You may also continue vitamins, steroids, antibiotics, water pills (diuretics). You may take over-the-counter painkillers such as Tylenol, Advil, ibuprofen, aspirin, and acetaminophen etc.
- 5. We ask that you try **not to eat approximately two hours** prior to the test. Part of the test makes you briefly dizzy and will often cause nausea. If you need to eat in the two hours prior to testing, please keep the meal light. Keep liquids to a minimum as well.



# Abraham Jacob, M.D. Department of Surgery/Division of Otolaryngology Otology, Neurotology, and Cranial Base Surgery

1501 N. Campbell Ave., 5OPC

Tucson, AZ 85724 Office: (520) 694-1000 Fax: (520) 694-6101

- 6. Please **do not wear contact lenses**. ENG/VNG testing requires measurement of eye movements and contact lenses interfere with accurate recordings. If you wear your contacts to the appointment, please bring your lens case.
- 7. Please do not wear <u>ANY</u> eye make-up and make sure any residual eye make-up is completely removed. Also, we would prefer that you have little to no make-up on your face, so please wear a minimal amount of **moisturizer**, **cream**, **lotion**, **or foundation make-up**. Face powder is okay.
- 8. Some patients experience slightly increased symptoms of dizziness after testing and you may wish to have someone available to drive you home.

Please call us if you have any questions.

#### **GENERAL INFORMATION**

To schedule, reschedule, or cancel an appointment: (520) 694-1000

Please complete any enclosed forms and bring them to your appointment. Please have your forms filled out prior to arriving to your appointment.

Although University Medical Center is a participating provider for many major insurance plans, it is the patient's responsibility to arrange any physician referrals or authorizations that may be required by your insurance plan.